Checklist (Prior to Shipping)

Includes Blood & Urine Specimens

1. Tubes
oxed Patient's first and last name, date of birth, gender, and date of collection
are written on all tubes and vials
☐ All the tubes and vials are capped tightly
2. Frozen
☐ Clear cap plastic vial (urine)
☐ 3 red top amber transfer tubes
☐ Lavender top clear transfer tube
☐ 3 ice packets
3. Room Temperature
☐ Royal-blue top Na-EDTA tube, trace mineral free
4. Test Requisition Form with Payment
☐ Test Requisition Form is complete
☐ Personal Health Assessment Form is complete
☐ Payment is included



LONDON · ASHEVILLE · ATLANTA

46 - 50 Coombe Road • New Malden, Surrey KT3 4QF • 020 8336 7750 • www.gdx.net/uk

©2012 Metametrix, Inc. All rights reserved IS-MM-10820 rev0317

IONSM Profile - Clinician

Blood Specimen Collection Instructions

This specimen collection kit can be used for the following test(s):

0090 ION Profile - Blood/Urine

0490 ION Profile w/Amino Acids 40 - Blood/Urine

Please Note: The ION Profile requires the patient to collect urine at home. This should be done prior to the blood collection. **All specimens, urine and blood, must be shipped together.** Patient must be fasting for blood draw. (Urine collection instructions are explained in the ION Profile - Clinician Specimen Collection Instructions.)

IMPORTANT:

All patient specimens require two unique identifiers **patient's name and date of birth**, as well as **date of collection**.

Patient's first and last name, date of birth, gender, and date of collection must be recorded on the **Test Requisition Form** as well as on all tube(s) and/or vial(s), using a permanent marker, or the test may not be processed.

Specimen

Serum, $^{\sim}9$ ml (3 tubes, $^{\sim}3$ ml each), frozen; **Plasma**, 2.5 - 3 ml, frozen;

Whole Blood, room temperature; Overnight Urine, 12 ml, frozen

Collection Materials

- 3 red/gray top serum separator tubes
- Royal-blue top Na-EDTA tube, trace mineral free
- Lavender top EDTA tube
- 3 red top amber transfer tubes
- Lavender top clear transfer tube
- 3 disposable pipettes

Shipping Materials

- Plastic shell tube tray
- 2 absorbent pads
- 3 ice packets
- Test Requisition Form
- Personal Health Assessment Form
- Biohazard bag with side pocket
- Specimen collection kit box

Please read all instructions carefully before you begin.

Patient Preparation

- It is best to ship the specimen within 48 hours of collection. Please refer to the
 enclosed shipping instructions before you collect to determine the days that the
 specimen can be shipped.
- Please check to make sure the patient has fasted prior to drawing blood.

Blood Collection

- 1. Write patient's first and last name, date of birth, gender, and date of collection on the Test Requisition Form (located in the pouch on top of the Specimen Collection Kit Box), as well as on all tube(s) and/or vial(s), using a permanent marker.
 - **IMPORTANT**: To ensure accurate test results, you **MUST** provide the requested information.
- 2. Freeze the ice packets.
- 3. Red/gray top serum separator tubes and red top amber transfer tubes
 - Draw the 3 red/gray top serum separator tubes .
 - Place upright in a rack at room temperature for 20 to 30 minutes to clot blood.
 - **Centrifuge** the red/gray top serum separator tubes for 15 minutes. The serum must be free of hemolysis or red blood cells.
 - **Pipette** all of the serum off of the 3 red/gray separator tubes into the 3 red top amber transfer tubes. (**DO NOT** fill more than ¾ full to allow for freezing). **Cap** each tube tightly.
 - Freeze the red top amber transfer tubes..
- 4. Royal-blue top Na-EDTA tube, trace mineral free
 - **Draw** the royal-blue top Na-EDTA tube completely.
 - Invert the tube gently 15 times immediately after the blood draw.
 - Leave the tube at room temperature. Do not centrifuge or transfer.
- 5. Lavender top EDTA tube and lavender top clear transfer tube
 - » **Draw** the lavender top EDTA tube completely.
 - » Invert the lavender top EDTA tube 10 times to mix the EDTA with the blood.
 - » **Centrifuge** immediately for 15 minutes. The plasma must be free of hemolysis or red blood cells.
 - » Remove the lavender top EDTA tube after centrifuging; DO NOT INVERT THE TUBE.
 - » **Pipette** plasma, using a fresh disposable pipette, 2.5 3 ml to the lavender top clear transfer tube.
 - » Freeze the lavender top clear transfer tube.

Specimen Preparation

- 1. Place all of the frozen transfer tubes and the frozen urine collection into the slots or the ends of the plastic shell tube tray (an exact fit is not necessary). Place the absorbent pad over the tubes. Place the frozen ice packets at each end of the tubes in the tray and one in the middle. Snap the tray closed. (Do not place the royal-blue top Na-EDTA tube inside the tray).
- 2. Place the tray, along with the royal-blue top Na-EDTA tube, into the biohazard bag.
- **3. Staple** payment to the bottom right-hand corner of the completed Test Requisition Form and **complete** the Personal Health Assessment Form; **Fold and Place** them in the side pocket of the biohazard bag.
- Seal the biohazard bag; Place it into the specimen collection kit box, and close the box.