



Requisition #: 9900001
Patient Name: Report Sample
Patient Age: 17
Patient Sex: M

Practitioner: NO PHYSICIAN
Date of Collection: 12/01/2022
Time of Collection: Not Given
Print Date: 03/21/2023
Report Date: 12/01/2021



Microbial Organic Acids Test

Metabolic Markers in Urine Reference Range (mmol/mol creatinine) Patient Value Reference Population - Males Age 13 and Over

Intestinal Microbial Overgrowth

Yeast and Fungal Markers

Table with 5 columns: Marker Name, Reference Range, Patient Value, and a graphical bar chart showing the patient's value relative to the reference range. Markers include Citramalic, 5-Hydroxymethyl-2-furoic, 3-Oxoglutaric, Furan-2,5-dicarboxylic, Furancarboxylglycine, Tartaric, Arabinose, Carboxycitric, and Tricarballic.

Bacterial Markers

Table with 5 columns: Marker Name, Reference Range, Patient Value, and a graphical bar chart. Markers include Hippuric, 2-Hydroxyphenylacetic, 4-Hydroxybenzoic, 4-Hydroxyhippuric, and DHPA (Beneficial Bacteria).

Clostridia Bacterial Markers

Table with 5 columns: Marker Name, Reference Range, Patient Value, and a graphical bar chart. Markers include 4-Hydroxyphenylacetic (C. difficile, C. stricklandii, C. lituseburens), HPPA (C. sporogenes, C. caloritolerans, C. botulinum), 4-Cresol (C. difficile), and 3-Indoleacetic (C. stricklandii, C. lituseburens, C. subterminale).

Additional Indicators

Table with 5 columns: Marker Name, Reference Range, Patient Value, and a graphical bar chart. Markers include 3-Hydroxy-3-methylglutaric and 2-Hydroxyhippuric.



Requisition #: 9900001
 Patient Name: Report Sample

Practitioner: NO PHYSICIAN
 Date of Collection: 12/01/2022

Indicator of Fluid Intake

21 *Creatinine 100 mg/dL

*The creatinine test is performed to adjust metabolic marker results for differences in fluid intake. Urinary creatinine has limited diagnostic value due to variability as a result of recent fluid intake. Samples are rejected if creatinine is below 20 mg/dL unless the client requests results knowing of our rejection criteria.

Explanation of Report Format

The reference ranges for organic acids were established using samples collected from typical individuals of all ages with no known physiological or psychological disorders. The ranges were determined by calculating the mean and standard deviation (SD) and are defined as $\pm 2SD$ of the mean. Reference ranges are age and gender specific, consisting of Male Adult (≥ 13 years), Female Adult (≥ 13 years), Male Child (< 13 years), and Female Child (< 13 years).


There are two types of graphical representations of patient values found in the new report format of both the standard Organic Acids Test and the Microbial Organic Acids Test.

The first graph will occur when the value of the patient is within the reference (normal) range, defined as the mean plus or minus two standard deviations.


The second graph will occur when the value of the patient exceeds the upper limit of normal. In such cases, the graphical reference range is "shrunk" so that the degree of abnormality can be appreciated at a glance. In this case, the lower limits of normal are not shown, only the upper limit of normal is shown.

In both cases, the value of the patient is given to the left of the graph and is repeated on the graph inside a diamond. If the value is within the normal range, the diamond will be outlined in black. If the value is high or low, the diamond will be outlined in red.

Example of Value Within Reference Range



Example of Elevated Value



Requisition #:	9900001	Practitioner:	NO PHYSICIAN
Patient Name:	Report Sample	Date of Collection:	12/01/2022

Interpretation

High tricarballic acid (propane-1,2,3-tricarboxylic acid) (Marker 9) could be caused by the intake of corn or corn-based food contaminated with fumonisins, a group of mycotoxins produced primarily by *F. verticillioides*, and other related species. Tricarballic acid is released from fumonisins during passage through the gastrointestinal tract. Tricarballic acid is an inhibitor of the enzyme aconitase and therefore interferes with the Krebs cycle. The main symptoms of aconitase deficiency are myopathy and exercise intolerance. It may also act as a magnesium chelator. Tricarballic acid is also metabolite of a component of a substance in modified corn starch, octenylsuccinic acid, found in a number of infant formulas such as Nutramigen, Vivonex, and Pregestimil. In addition, tricarballic acid is a byproduct of beet sugar and maple sugar refining and might appear after ingestion of these sugars. Tricarballic acid is also released from fumonisins upon certain food processing conditions. Clinical syndromes due to the intact mycotoxin are rare and characterized by abdominal pain and diarrhea. A specific role for fumonisins in the development of neural tube defects was suggested after the appearance of a cluster of such defects in Texas associated with consumption of corn from the heavily fumonisin-contaminated 1989 corn crop. More recent studies have shown that fumonisin B1 inhibits folate metabolism in cultured cells.

High 2-hydroxyphenylacetic acid (Marker 11) is associated with intestinal bacteria overgrowth and with the genetic disease phenylketonuria (PKU).

High 4-hydroxybenzoic acid and/or 4-hydroxyhippuric acid (Markers 12,13) may be due to bacterial overgrowth of the GI tract, intake of fruits such as blueberries rich in polyphenols (anthocyanins, flavonols, and hydroxycinnamates), or may be from paraben additive exposure. Parabens are 4-hydroxybenzoic acid alkyl esters with antimicrobial properties. 4-Hydroxybenzoic acid may be excreted as its glycine conjugate 4-hydroxyhippuric acid. High levels of these paraben metabolites in urine (>10 mmol/mol creatinine) may result from excessive exposure to parabens. Parabens are common preservatives allowed in foods, drugs, cosmetics and toiletries, but they also have a long history of use in a variety of pharmaceutical products for injection, inhalation, oral, topical, rectal or vaginal administration. Some individuals experience skin reactions as most parabens are readily and completely absorbed through the skin and the GI tract. Parabens have been considered safe because of their low toxicity profile and their long history of safe use; however, recent studies challenge this view. In 1998, Routledge *et.al.*, (Toxicol.Appl.Pharmacol. **153**,12-19), reported parabens having estrogenic activity *in vitro*. A number of *in vivo* studies have further elucidated potential endocrine disruption by parabens affecting reproduction or promote tumor growth. Parabens have been found at high levels in breast cancer biopsies, although a definitive relationship with breast cancer has not been demonstrated. Parabens may contribute to mitochondrial failure by uncoupling oxidative phosphorylation and depleting cellular ATP. 4-Hydroxyhippuric acid has been found to be an inhibitor of Ca²⁺-ATPase in end-stage renal failure. Eliminate all sources of parabens. To accelerate paraben excretion, use sauna therapy, the Hubbard detoxification protocol employing niacin supplementation, or glutathione supplementation (oral, intravenous, transdermal, or precursors such as N-acetyl cysteine [NAC]).

High DHPPA (3,4 dihydroxyphenylpropionic acid) (Marker 14) indicates excessive intake of chlorogenic acid, a common substance found in beverages and in many fruits and vegetables, including apples, pears, tea, coffee, sunflower seeds, carrots, blueberries, cherries, potatoes, tomatoes, eggplant, sweet potatoes, and peaches. Harmless or beneficial bacteria such as Lactobacilli, Bifidobacteria, and *E. coli* mediate the breakdown of chlorogenic acid to 3,4-dihydroxyphenylpropionic acid (DHPPA), and high values may indicate increased amounts of these species in the GI tract. In addition, one *Clostridia* species, *C. orbiscindens*, can convert the flavanoids luteolin and eriodictyol, occurring only in a relatively small food group that includes parsley, thyme, celery, and sweet red pepper to 3,4-dihydroxyphenylpropionic acid. The quantity of *Clostridia orbiscindens* in the GI tract is negligible (approximately 0.1% of the total bacteria) compared to the predominant flora of *Lactobacilli*, *Bifidobacteria*, and *E. coli*. Consequently, this marker is essentially useless as a general *Clostridia* marker but may be a good indicator of the presence of beneficial flora.

High 2-hydroxyhippuric acid (Marker 20) may result after ingestion of aspartame (NutraSweet®) or salicylates (aspirin), or from GI bacteria converting tyrosine or phenylalanine to salicylic acid. 2-Hydroxyhippuric acid is a conjugate of hydroxybenzoic acid (salicylic acid) and glycine.

<i>Requisition #:</i>	9900001	<i>Practitioner:</i>	NO PHYSICIAN
<i>Patient Name:</i>	Report Sample	<i>Date of Collection:</i>	12/01/2022

High quality nutritional supplements can be purchased through your practitioner or at New Beginnings Nutritionals, www.NBNUS.com <<http://www.NBNUS.com>> , or call 877-575-2467.

The nutritional recommendations in this test are not approved by the US FDA. Supplement recommendations are not intended to treat, cure, or prevent any disease and do not take the place of medical advice or treatment from a healthcare professional.